

Breast Cancer Research and Treatment

An open label, randomized phase 2 trial assessing the impact of food on the tolerability of abemaciclib in patients with Advanced Breast Cancer

--Manuscript Draft--

Manuscript Number:	BREA-D-22-00266R1
Full Title:	An open label, randomized phase 2 trial assessing the impact of food on the tolerability of abemaciclib in patients with Advanced Breast Cancer
Article Type:	Clinical trial
Keywords:	CDK4/6 inhibitor; Tolerability; breast cancer
Corresponding Author:	Elgene Lim, MBBS, FRACP, PhD Garvan Institute of Medical Research Darlinghurst, NSW AUSTRALIA
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	Garvan Institute of Medical Research
Corresponding Author's Secondary Institution:	
First Author:	Elgene Lim, MBBS, FRACP, PhD
First Author Secondary Information:	
Order of Authors:	Elgene Lim, MBBS, FRACP, PhD Francis Boyle Meena Okera Sherene Loi Sema Sezgin Goksu Gertjan van Hal Sonya C Chapman Jonathan Gable Yanyun Chen Gregory Price Anwar Hossein Corona Gainford Meritxell Bellet Ezquerra
Order of Authors Secondary Information:	
Abstract:	<p>Purpose: Abemaciclib, a CDK4 & 6 inhibitor, is indicated for advanced breast cancer treatment. Diarrhea is a frequently associated adverse event of abemaciclib. The study objective was to investigate if food intake impacts local gastrointestinal toxicity.</p> <p>Methods: This Phase 2 study (13Y-MC-JPCP, NCT03703466) randomized 72 patients 1:1:1 to receive abemaciclib 200mg monotherapy twice daily (1) with a meal, (2) in a modified fasting state or (3) without regard to food. Primary endpoints included: incidence of investigator assessed severe (\geqGrade 3), prolonged ($>$7-days) Grade 2 diarrhea, treatment discontinuation, dose modifications, and loperamide utilization during the first 3 cycles of treatment. Patient outcomes were captured via a daily electronic diary. Pharmacokinetics (PK) are reported.</p> <p>Results: Incidence of investigator assessed severe diarrhea (Grade \geq3) was 1.4% (1 patient in Arm 1). Median duration of Grade 3 diarrhea was 1 day by both investigator assessment (1 patient in Arm 1) and patient-reported assessment (1 patient each in</p>

	<p>Arms 1 and 3). Median duration of investigator-assessed Grade 2 diarrhea was 2 days overall. No patient discontinued treatment due to diarrhea. Nine patients (12.7%) had a dose reduction, and 7 patients (9.9%) had a dose omission due to diarrhea. Ninety-four percent of patients used loperamide at least once. Abemaciclib PK was comparable across the 3 arms.</p> <p>Conclusion: The results suggest that diarrhea incidence associated with abemaciclib was unrelated to timing of food intake, was predominantly low grade, of short duration and well managed with loperamide and dose modifications.</p>
<p>Response to Reviewers:</p>	<p>10th Jun 2022</p> <p>Dear reviewers and editor of Breast Cancer Research and Treatment,</p> <p>Thank you for considering this revised version of our manuscript (BREA-D-22-00266), titled “An open label, randomized phase 2 trial assessing the impact of food on the tolerability of abemaciclib in patients with Advanced Breast Cancer”.</p> <p>We have responded to each of the reviewer’s comments below and have provided a clean version of our resubmission with changes in response to reviewer comments indicated by page number and a document with all changes included as track changes. Thank you for the thorough review, we truly appreciate the attention and detailed comments that have greatly improved our submission.</p> <p>Yours sincerely,</p> <p>Prof. Elgene Lim (on behalf of all co-authors)</p> <p>Reviewer comments Reviewer #1</p> <p>1. Address potential reasons that this patient population had not yet received a CDK 4/6 inhibitor despite Palbociclib having been approved since 2015. Response: Palbociclib was approved in the US in early 2015 but our study was conducted in Europe and Australia. Palbociclib wasn’t approved until late 2016 in the EMA area and until mid-2017 in Australia. Therefore, palbociclib was not widely used in the study countries at the time of the clinical trial.</p> <p>2. Please address possible explanations that 41.7% of patients discontinued treatment, primarily due to progressive disease which is much higher than the reported rate of progressive disease in MONARCH 1 which was 25%. Response: We have clarified further in manuscript regarding the % of patients who discontinued treatment due to progressive disease. 21 patients (29.2%) discontinued treatment due to progressive disease in this study, which is slightly increased compared to the 25% in MONARCH 1. (Page 8)</p> <p>3. Revise manuscript to include more specific inclusion/exclusion criteria regarding the patients’ concomitant GI disorders as these disorders could confound the results. Response: We have expanded the exclusion criteria in the methods section to more comprehensively describe what patients were excluded from the study (Page 6). Patients were excluded if they had a serious concomitant systemic disorder (for example, active infection or a GI disorder causing clinically significant symptoms such as nausea, vomiting or diarrhea [such as Crohn’s disease, ulcerative colitis], or profound immune suppression) or a serious preexisting medical condition (for example, history of major surgical resection involving the stomach or small bowel) that, in the opinion of the investigator, would compromise/preclude the patient’s ability to adhere to the protocol.</p> <p>4. Address the limitation that there was no standardization of food intake, both regarding food composition and timing. Address how patients were monitored and confirmed to be taking the medication with food or without food according to the assigned randomization arm. Need clarification of the methods for validation of adherence within the randomization</p>

arms. If it can't be shown that there was control regarding the timing of the medications in regard to food, then how can we make any conclusions among the three arms?

Response: We have acknowledged that the composition of food intake across groups is a limitation in the manuscript now (Pge 12). The patient ediaries (Suppl Fig. 1) were used to monitor and confirm that patients were taking medication with or without food, the timing of medication intake with regard to food and the intake of food in the morning and evening. Patients completed ediaries daily, thus the short recall period would ensure accuracy in relation to confirming the taking and timing of medication with regard to food.

5. Please revise manuscript to include the investigators' role and responsibility in regard to the e diary. Were they supposed to review it and use that information in their investigator assessment of diarrhea? This needs to be clarified given that remarks are made in the limitations regarding the number of physicians who did or did not review the e diary. Were providers supposed to use PROs in treatment decisions or were to be blinded to PROs.

Response: Investigators had real time access to the e-diary data to facilitate monitoring, ensure timely patient education, and for the grading of diarrhea, but investigators were not required to review the e-diary data in their assessment of diarrhea as this information would be obtained through patient history during study visits. We discuss this in the statistical analysis section of the methods (Page 7).

Reviewer #2 - Although there are no novel findings from the study, it is an important study to conduct, and the paper is generally well written.

1. In the abstract, the primary objectives are clearly stated but the results make no reference to the primary endpoints.

Response: We have expanded the abstract to mention all primary endpoints in this section including incidence of investigator assessed severe (\geq Grade 3) diarrhea, treatment discontinuation due to diarrhea and dose modifications due to diarrhea.

2. Was the loperamide given prophylactically?

Response: Patients were instructed to commence over-the-counter antidiarrheal medication such as loperamide at the first sign of loose stools and antidiarrheal medication was not to be used prophylactically. We have now added this sentence for further clarity in page 6.

3. Fig 4: The labels/legend are not clear.

Response: We have updated the figure with better resolution.

Reviewer #3 - This randomized phase II trial used ePRO monitoring of diarrhea rates and patterns in MBC patients treated with abemaciclib in the fed vs modified-fasting state.

1. Was any guidance regarding diet given to patients to help them decrease abemaciclib- associated diet such as eating lower fat and lower fibre diets?

Response: No guidance was given to patients regarding diet apart from avoiding the consumption of grapefruit or grapefruit juice and other inducers and inhibitors of cytochrome P450 (CYP) 3A.

2. Were data collected on the diets that patients consumed?

Response: No data was collected on the diets that patients consumed.

3. Please provide data on the duration of grade 1 diarrhea by cycle for each of the arms.

Response: We do not have duration of Grade 1 diarrhea by cycle for each arm. We have included duration of Grade 1, 2 and 3 diarrhea by treatment arm across all cycles combined in Table 1.

4. A table or figure summarizing incidence and duration by cycle of grade 1 vs 2 vs 3 vs 4 diarrhea would be helpful.

Response: We do not have duration by cycle of grade 1 vs 2 vs 3. We have provided the incidence of diarrhea by cycle of grade 1 vs 2 vs 3 (there was no grade 4 diarrhea events noted in this study) in Fig. 4.

Reviewer #4 - The authors evaluated the effect of food intake on the tolerability of abemaciclib and reported that the incidence of diarrhea was not related to the timing of food intake. This study is compatible with the current interest of how to reduce the abemaciclib-related toxicity in the clinical practice since the cause of dose reduction or schedule modification are due to diarrhea.

1. The authors cited previous paper that a high-fat and high calorie meal modified the PK, and the NSAIDs paper that ingestion of food with NSAIDs reduced the GI adverse event. Based on those reports, what was the hypothesis? The food intake would reduce the GI event of abemaciclib?

Response: The study was a post-marketing commitment to the FDA. No hypothesis tests were planned for this study. Sample size was based on regulatory guidance (FDA). The references were mentioned to demonstrate that the intake of food can influence the PK and GI events like diarrhea.

2. Any previous reports regarding the food intake and oral chemotherapy toxicity?

Response: To our knowledge there are no previous reports examining food intake and oral chemotherapy toxicity.

3. Since abemaciclib is metabolized by CYP450, did the protocol specified the intake of grapefruits or any other CYP450 modulators?

Response: Patients were advised to avoid the consumption of grapefruit or grapefruit juice and other inducers and inhibitors of cytochrome P450 (CYP) 3A where possible. We have now included this in the manuscript (Page 6).

4. As the authors mentioned in the limitation section, the biggest concern about this study is the usage of loperamide. It is highly possible that loperamide masked the actual effect of food intake since the difference between three group would be small.

Response: We acknowledge that this is a limitation of the study and understand that this may have possibly masked the effect of food intake. Patients received diarrhea management guidance per label and were instructed to commence over-the-counter antidiarrheal medication such as loperamide at the first sign of loose stools, and not to be used prophylactically. Importantly, loperamide use was similar across arms, allowing us to examine the effect of food intake across arms in this study.

5. The sample size is too small to obtain the definitive conclusion.

Response: The sample size was based on regulatory guidance, not statistical powering (Page 7). We have revised our conclusion in the abstract and conclusion to be less definitive.

6. Since many patients have previously received many lines of chemotherapy, the nutrition status needs to be described (at least BMI).

Response: We have now included BMI data in Table 1.

7. Any washout period for the previous treatment was specified in the protocol?

Response: Patients were required to have discontinued all previous treatments for cancer and recovered from the acute effects of therapy. Per protocol, length of time between end of previous treatment and first abemaciclib dose was 14 to 28 days, depending on type of prior treatment. We have now included this in the manuscript

(Page 5).

Funding Information:

Eli Lilly and Company

Prof Elgene Lim

[Click here to view linked References](#)

1
2
3
4 **An open label, randomized phase 2 trial assessing the impact of food on the tolerability of**
5
6 **abemaciclib in patients with Advanced Breast Cancer**
7
8
9

10
11 **Authors:** Elgene Lim^{1#}, Frances Boyle^{2#}, Meena Okera³, Sherene Loi⁴, Sema Sezgin Goksu⁵, Gertjan
12 van Hal⁶, Sonya C. Chapman⁷, Jonathon Colby Gable⁸, Yanyun Chen⁸, Gregory L. Price⁸, Anwar M.
13 Hossain⁸, M. Corona Gainford⁸, Meritxell Bellet Ezquerro⁹
14
15
16
17

18 **Author Affiliations:** ¹Garvan Institute of Medical Research, St. Vincent's Clinical School, UNSW, NSW,
19 Australia; ²Mater Hospital, North Sydney, NSW, Australia; ³Adelaide Cancer Centre, Kurralta Park, SA,
20 Australia; ⁴Peter MacCallum Cancer Centre, Melbourne, VI, Australia; ⁵Akdeniz University Medical
21 Faculty, Antalya, Turkey; ⁶Eli Lilly and Company, Utrecht, The Netherlands; ⁷Eli Lilly and Company,
22 Windlesham, Surrey, UK; ⁸Eli Lilly and Company, Indianapolis, IN, USA; ⁹Hospital Universitario Vall
23 d'Hebron and Vall d'Hebron Institute of Oncology (VHIO), Barcelona, Spain.
24
25
26
27
28
29
30

31 # Contributed equally to the study
32
33

34 *** Corresponding Author:**
35

36 Elgene Lim
37
38

39 St. Vincent's Hospital Sydney,
40
41

42 Kinghorn Cancer Centre, 370 Victoria Street,
43
44

45 Darlinghurst, NSW, Australia 2010
46
47

48 e.lim@garvan.org.au
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 **ABSTRACT**
5
6

7 Purpose: Abemaciclib, a CDK4 & 6 inhibitor, is indicated for advanced breast cancer treatment. Diarrhea
8 is a frequently associated adverse event of abemaciclib. The study objective was to investigate if food
9 intake impacts local gastrointestinal toxicity.
10

11
12
13 Methods: This Phase 2 study (I3Y-MC-JPCP, NCT03703466) randomized 72 patients 1:1:1 to receive
14 abemaciclib 200mg monotherapy twice daily (1) with a meal, (2) in a modified fasting state or (3) without
15 regard to food. Primary endpoints included: incidence of investigator assessed severe (\geq Grade 3),
16 prolonged (>7 -days) Grade 2 diarrhea, treatment discontinuation, dose modifications, and loperamide
17 utilization during the first 3 cycles of treatment. Patient outcomes were captured via a daily electronic
18 diary. Pharmacokinetics (PK) are reported.
19
20
21
22
23
24
25

26 Results: Incidence of investigator assessed severe diarrhea (Grade ≥ 3) was 1.4% (1 patient in Arm 1).
27
28 Median duration of Grade 3 diarrhea was 1 day by both investigator assessment (1 patient in Arm 1) and
29 patient-reported assessment (1 patient each in Arms 1 and 3). Median duration of investigator-assessed
30 Grade 2 diarrhea was 2 days overall. No patient discontinued treatment due to diarrhea. Nine patients
31 (12.7%) had a dose reduction, and 7 patients (9.9%) had a dose omission due to diarrhea. Ninety-four
32 percent of patients used loperamide at least once. Abemaciclib PK was comparable across the 3 arms.
33
34
35
36
37
38

39 Conclusion: The results suggest that diarrhea incidence associated with abemaciclib was unrelated to
40 timing of food intake, was predominantly low grade, of short duration and well managed with loperamide
41 and dose modifications.
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

INTRODUCTION

Abemaciclib is an oral, selective, and potent inhibitor of cyclin-dependent kinases 4 and 6 (CDK4 & 6) dosed twice daily (BID) on a continuous schedule [1]. Resulting from the MONARCH series of clinical trials, abemaciclib is approved as monotherapy and in combination with endocrine therapy (ET) for the treatment of patients with hormone receptor-positive (HR+), human epidermal growth factor receptor 2-negative (HER2-) advanced breast cancer (ABC) [2].

MONARCH 1, a single-arm Phase 2 study of abemaciclib 200mg BID monotherapy in patients with refractory HR+, HER2- ABC, demonstrated promising clinical activity (objective response rate (ORR) of 19.7% (95% confidence interval [CI]: 13.3-27.5) [3].

MONARCH 2 was a randomized, double-blind, Phase 3 study of abemaciclib 150mg BID in combination with fulvestrant compared to placebo plus fulvestrant in women with HR+, HER2- ABC who had progressed following ET therapy [1]. This trial demonstrated significantly improved progression-free survival (PFS) (median 16.4 versus 9.3 months; hazard ratio [HR] [95% CI]: 0.553 [0.449-0.681]; $p < .001$) and overall survival (OS) (median 46.7 versus 37.3 months; HR [95% CI]: 0.757 [0.606-0.945]; $p = .01$) [1, 4].

MONARCH 3 was a randomized, double-blind Phase 3 study of abemaciclib 150mg BID in combination with a non-steroidal aromatase inhibitor (NSAI) compared to placebo plus NSAI as initial therapy in women with HR+, HER2- ABC [5]. Abemaciclib plus NSAI significantly improved PFS (median 28.18 versus 14.76 months; HR [95% CI]: 0.540 [0.418-0.698]; $p = .000002$).

Diarrhea was the most frequently reported treatment-emergent adverse event (TEAE) of any-grade in patients prescribed abemaciclib in all three pivotal studies, irrespective of whether abemaciclib was taken as a monotherapy (MONARCH 1 [200mg BID]: diarrhea=90.2%) or in combination with ET (MONARCH 2 and MONARCH 3 [150mg BID]: diarrhea=87.1% and 82.3% respectively) [1, 3, 5]. Grade 3 diarrhea was reported in 20% of patients in MONARCH 1, 14% in MONARCH 2 and 10% in MONARCH 3 [3-5]. In all three trials, incidence of Grade 2 and Grade 3 diarrhea was greatest during the first month of treatment and decreased over the remaining cycles of therapy (Fig. 1). The median duration of any-grade diarrhea

1
2
3
4 was similar across the 3 trials ranging 6-8 days, with the median duration of Grade 2 diarrhea ranging 8-
5
6 11 days and Grade 3 diarrhea, 5-8 days [1, 3, 5]. In each study, diarrhea was retrospectively assessed by
7
8 the investigator at the beginning of each 28-day cycle and graded as per Common Terminology Criteria
9
10 for Adverse events (CTCAE) criteria [1, 3, 5]. To our knowledge, no patient-reported daily data on
11
12 abemaciclib and diarrhea have been published to date.

13
14
15 Standardized anti-diarrheal management plans are outlined in the product label [2]. Patients are advised
16
17 to commence over-the-counter (OTC) antidiarrheal medication such as loperamide at the first sign of
18
19 loose stools, increase their fluid intake, and contact their physician. If diarrhea has not resolved within 24
20
21 hours, abemaciclib should be suspended until resolution occurs. Resolution is defined as either a
22
23 reduction to baseline or Grade 1 (<4 stools per day increase over baseline). The label also includes
24
25 detailed guidance for dose modifications and reductions according to severity of diarrhea [2].
26
27

28 Consistent with management guidance, 21%, 19%, and 14% of patients in MONARCH 1, 2, and 3
29
30 respectively required dose reductions and 1%, 3%, and 2% respectively discontinued the study drug due
31
32 to diarrhea [1, 3, 5]. Antidiarrheal use across the studies varied between 61% in MONARCH 1, 76% in
33
34 MONARCH 2, and 61% in MONARCH 3.
35

36
37 In MONARCH 2 and 3, a paper version of the patient-reported outcomes (PRO) assessment was
38
39 completed once at baseline, and on treatment; with more frequent collection in earlier cycles, and at the
40
41 follow-up visit [6, 7]. PRO results from abemaciclib in combination with fulvestrant (MONARCH 2) or in
42
43 combination with NSAID (MONARCH 3) did not show clinically significant differences in patient-reported
44
45 global health, functioning, or most symptoms compared to ET alone [6, 7]. Baseline scores were similar
46
47 between treatment arms in each study. In both trials, diarrhea was the only patient-reported symptom with
48
49 a statistically significant and clinically meaningful difference between treatment arms. These diarrhea
50
51 findings were reported in early treatment cycles, consistent with investigator assessments, decreased in
52
53 later cycles and returned to near baseline levels at the post-therapy follow-up visits [6, 7].
54

55
56 For some drugs, coadministration with food can impact bioavailability and may have clinically significant
57
58 consequences. In clinical studies, a high-fat, high-calorie meal increased the exposure (AUC) of
59
60 abemaciclib analytes by 9% and increased C_{max} by 25% [8]. These changes in exposure are not clinically
61

1
2
3
4 meaningful and abemaciclib is therefore given without regard to food. However, it is possible that taking
5
6 abemaciclib with food may impact local gastrointestinal toxicity independently of systemic
7
8 pharmacokinetics (PK) and thus alter drug tolerability. As an example, ingestion of food with nonsteroidal
9
10 anti-inflammatory drugs (NSAIDs) is often preferred because it reduces local gastrointestinal adverse
11
12 effects [9]. In order to address this issue and at the request of the U.S. Food and Drug Administration
13
14 (FDA), this randomized study (I3Y-MC-JPCP) evaluated the impact of coadministration of food on the
15
16 incidence and tolerability of diarrhea in patients with HR+, HER2- ABC receiving abemaciclib
17
18 monotherapy.
19
20
21
22

23 **METHODS**

24 *Study design and patients*

25
26
27 JPCP (NCT03703466) was a global, randomized, open-label Phase 2 study evaluating the timing of food
28
29 intake on the incidence of severe diarrhea (Grade ≥ 3) or prolonged Grade 2 diarrhea (>7 days duration)
30
31 when receiving abemaciclib monotherapy 200mg orally (PO) BID in patients with previously treated HR+,
32
33 HER2- ABC.
34
35
36

37
38 The study was conducted at 15 centers in 5 countries (Australia, Belgium, Russian Federation, Spain and
39
40 Turkey). It was approved by ethical and local institutional review boards for the participating sites and was
41
42 conducted according to the Declaration of Helsinki. Patients provided written informed consent prior to
43
44 trial enrollment. The study was overseen by an ethics review board.
45
46

47 Eligible patients included males and females ≥ 18 years of age with a diagnosis of recurrent, locally
48
49 advanced, unresectable, or metastatic HR+, HER2- ABC. Patients were required to have an Eastern
50
51 Cooperative Oncology Group (ECOG) performance status of ≤ 1 ; a willingness to use an e-diary; and
52
53 having no prior use of a CKD4 & 6 inhibitor. Patients must have received ≥ 1 chemotherapy regimen and
54
55 progressed after prior anti-estrogen therapy for ABC. Patients were required to have discontinued all
56
57 previous treatments for cancer and recovered from the acute effects of therapy. Per protocol, length of
58
59 time between end of previous treatment and first abemaciclib dose was 14 to 28 days, depending on type
60
61
62
63
64
65

1
2
3
4 of prior treatment. Patients were ineligible if they had a serious concomitant systemic disorder (for
5
6 example, active infection or a gastrointestinal disorder causing clinically significant symptoms such as
7
8 nausea, vomiting or diarrhea [such as Crohn's disease, ulcerative colitis], or profound immune
9
10 suppression) or a serious preexisting medical condition (for example, history of major surgical resection
11
12 involving the stomach or small bowel) that, in the opinion of the investigator, would compromise/preclude
13
14 the patient's ability to adhere to the protocol.
15

16
17 Patients were randomly assigned 1:1:1 to take abemaciclib 200mg monotherapy either; with a meal (Arm
18
19 1); in a modified fasting state defined as at least 1 hour before or 2 hours after a meal (Arm 2); or without
20
21 regard to food (Arm 3). A meal was defined as whatever the patient would normally eat at that time.
22

23 Patients were advised to avoid the consumption of grapefruit or grapefruit juice and other inducers and
24
25 inhibitors of cytochrome P450 (CYP) 3A where possible, as these can affect the exposure of abemaciclib.
26

27 Abemaciclib was administered on a continuous BID schedule with at least 6 hours separating doses.

28
29 Treatment cycles lasted 28 days. The study period consisted of the first 3 cycles of treatment for each
30
31 individual patient and all results presented herein reflect the first 3 cycles only. Patients who continued to
32
33 receive benefit following cycle 3 remained on treatment at investigator discretion and took abemaciclib
34
35 without regard to food as per label. All study procedures were followed until study completion, which
36
37 occurred when the last enrolled patient completed 3 cycles. Loperamide was the protocol specified anti-
38
39 diarrheal medication. Patients received diarrhea management guidance per label and were provided with
40
41 written support materials.
42

43
44 A training program using both face-to-face guidance and virtual media was developed to ensure
45
46 standardized implementation of the e-diary globally. All patients completed a daily e-diary detailing the
47
48 timing of each abemaciclib dose in relation to a meal, number of bowel movements (BM), and number of
49
50 loperamide tablets. Supplemental Fig. 1 visualizes the patient's e-diary experience and how they were
51
52 prompted to respond within the e-diary. To determine an accurate baseline assessment, patients
53
54 recorded daily number of BM for a week prior to study commencement. Adherence to study medication
55
56 and e-diary was made available for investigator and site staff review.
57
58
59
60
61
62
63
64
65

1
2
3
4 PK samples were collected prior to first dose on cycle 1 day 1 and then in conjunction with other
5
6 laboratory samples in cycle 1: day 15, cycle 2: days 1 and 15, and cycle 3: day 1.
7

8 9 *Objectives*

10
11 The primary objective was to evaluate and summarize investigator-assessed incidence of Grade ≥ 3 and
12 prolonged Grade 2 (>7 days continuous duration) diarrhea; dose reductions, interruptions, and
13 discontinuations due to diarrhea; and patient-reported utilization of anti-diarrheal medications during the
14 first 3 cycles of treatment. Secondary objectives included overall safety, incidence and severity of TEAEs,
15 serious adverse events, deaths and clinical laboratory abnormalities. PK analysis included steady-state
16 concentrations of abemaciclib, its active metabolites LSN2839567 (M2) and LSN3106726 (M20), and
17 total active analytes (sum of abemaciclib+M2+M20). An exploratory objective was to evaluate and
18 summarize incidence and duration of diarrhea reported daily by the patient using an e-diary.
19
20
21
22
23
24
25
26
27

28 29 *Statistical Analysis*

30
31 The study planned to enroll approximately 60 patients and was descriptive in nature. It was not powered
32 for formal statistical comparison between groups, and sample size was based on regulatory guidance.
33
34 Assignment to treatment arms were determined by a computer-generated random sequence and
35 randomization was stratified by region.
36
37
38
39

40 Baseline analyses and patient dispositions were based on the intent-to-treat (ITT) population which
41 included all patients enrolled and randomized in the trial. Safety analyses included all patients who
42 received at least 1 dose of abemaciclib. Analyses of all primary investigator-assessed endpoints and
43 adverse events were based on the safety population and summarized by study arms. Investigator
44 assessments of diarrhea and dose modifications were required on day 1 of each cycle and were
45 conducted as per their usual practice. While investigators and site staff had real time access to e-diary
46 data to facilitate monitoring, ensure timely patient education, and for the grading of diarrhea, investigators
47 were not required to review the e-diary data **in their assessment of diarrhea**. Adherence with study
48 medication and e-diary was monitored and recorded by investigators. Patient data collected via the e-
49 diary was summarized descriptively to provide daily estimates of loperamide use, compliance with
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 assignment to food administration cohort, as well as incidence and duration of diarrhea, complementary
5
6 to investigator assessment. PK analysis included all patients who had received at least 1 dose of
7
8 abemaciclib and who had at least 1 evaluable PK sample.
9

10 11 12 13 **RESULTS**

14 15 16 *Patients*

17
18
19 Seventy-two patients were enrolled from December 4th, 2018 to April 10th, 2019 with 24 patients
20
21 randomized to each of the 3 arms (Fig. 2). One patient in Arm 2 withdrew from the study and did not
22
23 receive treatment, resulting in a safety population of 71 patients. Only 1 patient was male; this patient was
24
25 randomized to Arm 1 (Table 1). The majority of patients (71%) were recruited from Australia, Spain, and
26
27 Belgium. The mean age (years) was similar in Arm 1 (58), Arm 2 (57), and Arm 3 (58). Approximately
28
29 22% of patients had received ≥ 5 prior chemotherapies for ABC. Mean treatment compliance calculated by
30
31 pill return across the 3 groups was 95.1%. Mean compliance with the assigned arm's timing of
32
33 abemaciclib dosing in relation to a meal was 97.4%. Mean e-diary completion compliance across all
34
35 study arms was 95.7%.
36

37
38 Although only 2 of 18 investigators assessed the e-diary portal 3 or more times during the study period,
39
40 the portal was accessed more frequently by study coordinators and nurses at each site. At the July 8th,
41
42 2019 data cutoff, 30 patients (41.7%) had discontinued treatment, with 21 patients (29.2%) discontinuing
43
44 due to progressive disease.
45

46 47 *Diarrhea incidence and dose modifications*

48
49 Food did not appear to impact incidence of investigator assessed diarrhea or dose modifications. While
50
51 the numbers were low, the incidence of Grade ≥ 3 and prolonged Grade 2 diarrhea during the first 3 cycles
52
53 of treatment were comparable across the 3 arms (Table 2). Notably, investigator assessed Grade 3
54
55 diarrhea occurred in only 1 patient overall (1.4%) during cycle 1 in Arm 1 and lasted 1 day. Prolonged
56
57 Grade 2 diarrhea (>7 -days) occurred in 11 patients overall (15.5%).
58
59
60
61
62
63
64
65

1
2
3
4 Dose reductions and omissions (dose interruptions) were balanced across Arms 1, 2, and 3 with 9
5
6 (12.7%) of all 71 patients having at least 1 dose reduction due to diarrhea and 7 (9.9%) of all patients
7
8 requiring a dose omission for diarrhea (Table 2). 94% of patients used loperamide at least once in the first
9
10 3 cycles. Dose reduction due to fatigue and neutropenia were also balanced across all 3 arms, occurring
11
12 in 4 (5.6%) and 8 (11.3%) of all 71 patients respectively.
13

14 *Pharmacokinetics*

15
16
17 A total of 69 patients had evaluable PK plasma samples: 24, 21, and 24 in Arms 1, 2, and 3 respectively.
18

19 A total of 260, 256, and 259 samples were available for abemaciclib, and its active metabolites M2 and
20
21 M20 respectively. Abemaciclib concentrations were comparable across the 3 treatment arms (Fig. 3), with
22
23 average values ranging from 305 to 369 ng/ml on cycle 1 day 15. The mean range for total active
24
25 analytes for the same time point was similarly comparable (1.38–1.60 uM).
26
27

28 *Treatment-Emergent Adverse Events (TEAEs)*

29
30
31 Investigator-reported TEAEs of diarrhea were experienced by 21 patients (87.5%) in Arm 1, 22 patients
32
33 (95.7%) in Arm 2, and 22 patients (91.7%) in Arm 3. No Grade 4 or 5 diarrhea events were reported
34
35 (Table 2). TEAEs were otherwise consistent with prior abemaciclib trials and are listed in Table 3. No
36
37 patients experienced a serious AE of diarrhea, or discontinued abemaciclib due to diarrhea, during the
38
39 first 3 cycles. Importantly, despite the frequent use of loperamide, no patient experienced Grade 3 or
40
41 greater constipation. Overall, 2 patients experienced Grade 1 constipation and 1 patient had a Grade 2
42
43 event. Two deaths occurred while on therapy or within 30 days of treatment discontinuation, both of which
44
45 were associated with the study disease and assessed by the investigators to be unrelated to the study
46
47 treatment. One death (4.2%) in Arm 1 was due to respiratory failure related to pulmonary progressive
48
49 disease and pulmonary infection, and 1 death (4.3%) in Arm 2 was due to abnormal hepatic function
50
51 where the patient had been diagnosed with liver metastasis prior to starting treatment. The most common
52
53 treatment-emergent laboratory toxicities experienced by all patients included increased creatinine and
54
55 decreased neutrophil count. Increases in serum creatinine levels are a known pharmacodynamic effect of
56
57 abemaciclib caused by inhibition of renal tubular secretion without affecting glomerular function [10]. The
58
59
60
61
62
63
64
65

1
2
3
4 TEAE of neutropenia has been observed across MONARCH studies. Neutropenia was not associated
5
6 with severe infection [1, 3, 5].
7

8
9 Diarrhea incidence and duration as assessed by the investigator (A and B) and the patient (C and D)
10 during the first 3 cycles of therapy is illustrated in Fig. 4. This visually highlights the discrepancy between
11 both assessments. One episode of Grade 3 diarrhea was reported by investigator assessment (for 1
12 patient in Arm 1) while 2 episodes were reported by patient assessment (1 patient each in Arm 1 and Arm
13 3). All reported Grade 3 episodes were of 1 day's duration. Grade 2 diarrhea as assessed by the
14
15 investigator was of longer duration than that assessed by the patient (median duration 2 days vs 1 day).
16
17 Median duration of investigator-assessed Grade 2 diarrhea was 2 days overall: 2 days (Arm 1), 7 days
18 (Arm 2), and 3 days (Arm 3). Finally, Fig. 4A demonstrates that Grade 1 diarrhea was assessed by the
19 investigator as continuous while patient daily assessment via the e-diary (Fig. 4C) reported intermittent
20 and short duration Grade 1.
21
22
23
24
25
26
27
28

29
30 Loperamide doses and corresponding patient-reported grade of diarrhea (calculated from daily numbers
31 of BM relative to baseline recorded in the patient e-diaries) are displayed in Fig. 4E. While loperamide
32 was used frequently, the number of tablets taken varied daily with the grade of diarrhea experienced.
33
34
35
36
37
38

39 DISCUSSION

40
41
42 Results from this trial demonstrate that food does not appear to impact the incidence or duration of
43 prolonged Grade 2 and Grade ≥ 3 diarrhea, supporting the product label that abemaciclib may be taken
44 with or without food. Abemaciclib PK was comparable across the 3 treatment arms, indicating no major
45 differences in drug exposure related to meal timing. It has been reported that a high-fat, high-calorie meal
46 increases abemaciclib plasma concentrations to a non-clinically meaningful extent [8]. However, the
47 conditions of this trial simulate real-world drug administration rather than the high-fat, high-calorie meals
48 in clinical studies specifically designed to assess the effect of food on PK parameters.
49
50
51
52
53
54
55

56
57 The findings reported here include several novel insights about abemaciclib-associated diarrhea beyond
58 evaluation of the effect of food. Electronic capture of patient-reported outcomes (ePRO) can provide more
59
60
61

1
2
3
4 granular reporting of adverse events than can be captured by traditional techniques [11]. Investigator
5 assessments are complemented by daily patient reports of BM frequency and antidiarrheal use recorded
6 with easy-to-use handheld e-diaries (Figs 4A and B). Capturing daily patient-reported data via e-diary
7 facilitated detailed, real-time characterization of the patient experience compared to retrospective
8 physician assessment alone. Mean compliance with daily completion of the e-diary was 96% and is
9 remarkable for a global trial in a patient population with heavily pre-treated ABC. This suggests that it is
10 feasible, with appropriate training and oversight, to employ such tools in oncology trials. The overall
11 incidence of diarrhea (any-grade) in the current study was similar to that in MONARCH 1, where the dose
12 used, and patient population were the same. However, the incidence of Grade 3 diarrhea was much lower
13 here (1% vs 20%). This could suggest that as physicians have gained more experience with abemaciclib
14 and incorporated standard diarrhea management guidelines into clinical care, abemaciclib-associated
15 diarrhea is better understood and better managed. An assessment of the baseline number of BM as
16 performed in this study would be important to allow an accurate grading of diarrhea as per CTCAE
17 criteria. The median duration of Grade 2 and Grade 3 diarrhea, reported by both investigators and
18 patients, was shorter than what was observed in the MONARCH 1, 2, and 3 trials [3-5]. Interestingly,
19 patients in this study reported earlier resolution of Grade 2 diarrhea than investigators (Fig. 4).
20 Furthermore, Fig. 4 illustrates that patients reported intermittent and shorter durations of Grade 1 diarrhea
21 as opposed to the continuous Grade 1 diarrhea assessed by investigators. These data demonstrate the
22 benefits of accurately recording grade and duration of diarrhea and reinforces the importance of recording
23 the patient experience in real time. The additional granularity obtained through use of the daily e-diary
24 permitted exploration of the relationship between antidiarrheal use and the incidence, grade, and duration
25 of diarrhea. While loperamide was frequently used over the course of the study (Fig. 4C), the pattern of
26 use suggests patients were appropriately educated regarding diarrhea management and varied their use
27 of loperamide according to the severity experienced.

28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54 Consistent with management guidelines in the product label allowing for dose modifications for diarrhea,
55 12.7% of patients in this study had at least 1 dose reduction. A previous exploratory analysis
56 demonstrated no difference in PFS outcomes for patients who dose reduced versus those who did not in
57 the MONARCH 1, 2, and 3 trials, suggesting patients can receive a reduced dose and still derive
58
59
60
61

1
2
3
4 meaningful benefit from abemaciclib treatment [12]. Those findings and the results of this study
5
6 collectively suggest the standardized diarrhea management plan described in the product label
7
8 appropriately manages abemaciclib-associated diarrhea without risking a decrease in efficacy.
9

10
11 This study had several limitations. First, it was of modest size limiting the ability to make comparisons
12
13 between arms. Additionally, while data collected via e-diary was made available to site staff in real-time,
14
15 e-diary user metrics reports showed it was predominantly accessed by study coordinators and nurses
16
17 with only 2 of 18 investigators accessing the data ≥ 3 times in the 3-month study period (data on file). Ten
18
19 of 18 investigators never accessed the e-diary data. This potentially explains the discrepancies seen
20
21 between investigator and patient assessments (Fig. 4). Although high compliance rates with the daily
22
23 patient e-diaries provided valuable insights into diarrhea incidence and management, it could be argued
24
25 that patients may have been biased towards better management because of increased awareness and a
26
27 daily requirement to enter data. Prior studies have indicated that active assessment and monitoring of
28
29 symptomatic adverse events could yield positive outcomes for patients, including health related quality of
30
31 life [13, 14]. Further studies are needed to understand the impact of patient self-monitoring on outcomes.
32
33 Loperamide use was higher than what was previously seen in the MONARCH studies and it is possible
34
35 that the high use of anti-diarrheal medication masked the impact of food on the incidence and severity of
36
37 diarrhea. Food composition intake was also not standardized across groups. Finally, while PK analysis
38
39 detected no discernable differences between the 3 arms, the study was not designed to assess the
40
41 impact of food on abemaciclib PK, and it is likely that only large effects would be detected with a non-
42
43 crossover study design and sparse PK sampling.
44

45
46 In conclusion, the results suggest that abemaciclib can be taken with or without food. Management
47
48 guidance for associated diarrhea provided in the product label is appropriate. The current study results
49
50 suggest that diarrhea is manageable with loperamide and dose modifications as needed. This is
51
52 demonstrated by a progressive decrease in the frequency of Grade 3 diarrhea (in the first 3 cycles) for
53
54 MONARCH 1, 2, and 3, and in the present study. Diarrhea onset is usually in the first cycle, and patient-
55
56 reported e-diary data in the present study shows shorter duration of Grade 1 and 2 diarrhea than reported
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

by investigator assessment. It is critical that patients receiving abemaciclib are educated on how to manage diarrhea and commence loperamide at the first sign of loose stools for optimal treatment benefit.

1
2
3
4 **DATA AVAILABILITY**
5

6
7 Eli Lilly and Company makes patient-level data available from Lilly-sponsored studies on marketed drugs
8
9 for approved uses following acceptance for publication. Lilly is one of several companies that provide this
10
11 access through the website clinicalstudydatarequest.com. Qualified researchers can submit research
12
13 proposals and request anonymized data to test new hypotheses. Lilly's data sharing policies are provided
14
15 on the clinicalstudydatarequest.com site under the Study Sponsors page.
16
17

18
19
20 **ACKNOWLEDGEMENTS**
21

22 The authors are grateful to the patients, their families, and caregivers for participating in study JPCP, and
23
24 to the study investigator and site staff for their collaboration. Writing and editing provided by Nicholas
25
26 Pulliam, Sandra Deady and John Hurley of Eli Lilly and Company. Funding provided by Eli Lilly and
27
28 Company.
29
30

31
32 **AUTHOR CONTRIBUTIONS**
33

34 All authors contributed to the intellectual content of the work through conception, design, data acquisition,
35
36 analysis and interpretation as well as the drafting and critical review of the manuscript. All authors
37
38 provided their final approval of the manuscript.
39
40
41

42 **COMPETING INTERESTS INFORMATION**
43

44
45 EL reports a consulting or advisory (paid to his institution) role with Eli Lilly and Company, Novartis
46
47 Australia, Pfizer and Roche; institutional research funding from Novartis Australia; travel or
48
49 accommodation expenses from Eli Lilly and Company; and a patent/royalty/intellectual property for the
50
51 sensitization of BCL-2-expressing breast tumors to chemotherapy by the BH3 mimetic ABT-737. FB
52
53 reports a consulting or advisory role with Pfizer, Eli Lilly and Company, Novartis Australia, Roche and
54
55 Eisai; and travel or accommodation expenses from Novartis Australia. MO has no competing interests to
56
57 declare. SL reports receiving research funding to her institution from Novartis, Bristol Meyers Squibb,
58
59 Merck, Puma Biotechnology, Eli Lilly, Nektar Therapeutics Astra Zeneca, Roche-Genentech and Seattle
60
61
62
63
64
65

1
2
3
4 Genetics; a consulting or advisory role (not compensated) with Seattle Genetics, Novartis, Bristol Meyers
5
6 Squibb, Merck, AstraZeneca and Roche-Genentech; a consulting or advisory role (paid to her institution)
7
8 with Aduro Biotech, Novartis, GlaxoSmithKline, Roche-Genentech, Astra Zeneca, Silverback
9
10 Therapeutics, G1 Therapeutics, PUMA Biotechnologies, Pfizer, Seattle Genetics and Bristol Meyers
11
12 Squibb; is a Scientific Advisory Board Member of Akamara Therapeutics; and is supported by the
13
14 National Breast Cancer Foundation of Australia Endowed Chair and the Breast Cancer Research
15
16 Foundation, New York. SSG reports a consulting or advisory role with Novartis, MSD, Roche and Amgen;
17
18 speakers' bureau company with Eli Lilly and Company, Novartis, Roche, Phizer and BMS; travel,
19
20 accomodation or congress fees from Pfizer, Novartis, Roche, Astra-Zeneca; and is a primary/sub
21
22 investigator of many clinical studies sponsored by Roche, MSD, BMS and Novartis. SCC, JCG, YC, GLP,
23
24 AMH and MCG are full-time employees of Eli Lilly and Company and are Eli Lilly and Company
25
26 shareholders. MB-E reports personal fees from Pfizer, Novartis, and Eli Lilly and Company; and travel
27
28 expenses from Pfizer and Roche.
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figure Legends

Fig. 1 Comparison of investigator-assessed treatment-emergent diarrhea in patients receiving abemaciclib from studies JPCP, MONARCH 1, MONARCH 2, and MONARCH 3

Fig. 2 Consort diagram

Fig. 3 Abemaciclib concentrations on Cycle 1 Day 15 across the three treatment arms

Fig. 4 **Duration and incidence of diarrhea in cycles 1-3**

- A. Investigator assessed
- B. **Investigator assessed**
- C. Patient-reported e-diary
- D. **Patient-reported e-diary**
- E. Patient-reported e-diary account of loperamide doses in cycles 1-3

Suppl Fig. 1 Patient facing e-diary

Table Legends

Table 1 Demographics and baseline characteristics

Table 2 Primary endpoint results, investigator assessed

Table 3 Treatment-emergent adverse events (TEAEs) \geq Grade 3; investigator-assessed

Table 1 Demographics and baseline characteristics

	Overall	Arm 1	Arm 2	Arm 3
	N = 72	N = 24	N = 24	N = 24
Sex, n (%)				
Female	71 (98.6)	23 (95.8)	24 (100.0)	24 (100.0)
Male	1 (1.4)	1 (4.2)	0	0
Age categories, n (%)				
<65 years	53 (73.6)	17 (70.8)	15 (62.5)	21 (87.5)
≥65 years	19 (26.4)	7 (29.2)	9 (37.5)	3 (12.5)
Age (years)				
Mean	57.6	58.2	56.9	57.6
Race, n (%)				
Asian	2 (2.8)	2 (8.3)	0	0
White	70 (97.2)	22 (91.7)	24 (100.0)	24 (100.0)
BMI, kg/m² (SD)				
Mean	26.9 (4.5)	26.6 (4.3)	28.3 (5.4)	26.0 (3.6)
Region, n (%)				
Turkey / Russia	21 (29.2)	7 (29.2)	7 (29.2)	7 (29.2)
Australia / Spain / Belgium	51 (70.8)	17 (70.8)	17 (70.8)	17 (70.8)
ECOG Performance Status, n (%)				
0	37 (51.4)	13 (54.2)	10 (41.7)	14 (58.3)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1	34 (47.2)	11 (45.8)	13 (54.2)	10 (41.7)
Nature of disease, n (%)				
Visceral	59 (81.9)	20 (83.3)	17 (70.8)	22 (91.7)
Bone only	8 (11.1)	3 (12.5)	5 (20.8)	0
Others	4 (5.6)	1 (4.2)	1 (4.2)	2 (8.3)
Missing	1 (1.4)	0	1 (4.2)	0
Number of organ sites (%)				
1	14 (19.4)	3 (12.5)	8 (33.3)	3 (12.5)
2	24 (33.3)	9 (37.5)	5 (20.8)	10 (41.7)
≥3	33 (45.8)	12 (50.0)	10 (41.7)	11 (45.8)
Number of prior chemotherapy regimens for ABC, n (%)				
<5	54 (75.0)	20 (83.3)	14 (60.9)	20 (83.3)
≥5	16 (22.2)	4 (16.7)	9 (39.1)	3 (12.5)

Intention-to-treat (ITT) population. All arms received 200mg abemaciclib monotherapy twice per day: Arm 1 = taken with a meal; Arm 2 = taken without a meal (modified fasting condition); Arm 3 = taken without regard to food. Abbreviations: ECOG = Eastern Cooperative Oncology Group; N = number of patients in the intent-to-treat population; n = number of patients within category. ECOG status missing for 1 patient in Arm 2

Table 2 Primary endpoint results, investigator assessed

	Overall	Arm 1	Arm 2	Arm 3
	N = 71	N = 24	N = 23*	N = 24
Endpoint				
Diarrhea (any grade), n (%)	65 (91.5)	21 (87.5)	22 (95.7)	22 (91.7)
Grade 1 diarrhea, n (%)	35 (49.3)	10 (41.7)	14 (60.9)	11 (45.8)
Duration of Grade 1 diarrhea, median days	8	9	6	10
Grade 2 diarrhea, n (%)	29 (40.8)	10 (41.7)	8 (34.8)	11 (45.8)
Duration of Grade 2 diarrhea, median days	2	2	7	3
Grade 2 diarrhea lasting >7 days, n (%)	11 (15.5)	2 (8.3)	4 (17.4)	5 (20.8)
Grade 3 diarrhea, n (%)	1 (1.4)	1 (4.2)	0	0
Duration of Grade 3 diarrhea, median days	1	1	0	0
Grade 4 diarrhea, n (%)	0	0	0	0
≥1 Dose reduction due to diarrhea, n (%)	9 (12.7)	4 (16.7)	2 (8.7)	3 (12.5)
≥1 Dose omission due to diarrhea, n (%)	7 (9.9)	4 (16.7)	1 (4.3)	2 (8.3)
Treatment discontinued due to diarrhea, n (%)	0	0	0	0

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Loperamide use, n (%)	67 (94.3)	23 (95.8)	21 (91.3)	23 (95.8)
-----------------------	-----------	-----------	-----------	-----------

All arms received 200mg abemaciclib monotherapy twice per day: Arm 1 = taken with a meal; Arm 2 = taken without a meal (modified fasting condition); Arm 3 = taken without regard to food. First three treatment cycles only. N = number of patients receiving at least 1 abemaciclib dose (safety population); n = number of patients within category

*one patient in Arm 2 discontinued from the study prior to receiving treatment

Table 3 Treatment-emergent adverse events (TEAEs) Grade \geq 3; investigator-assessed

	Overall	Arm 1	Arm 2	Arm 3
	N = 71	N = 24	N = 23	N = 24
TEAE	n (%)	n (%)	n (%)	n (%)
Neutropenia	20 (28.2)	7 (29.2)	3 (13.0)	10 (41.7)
Leukopenia	9 (12.7)	5 (20.8)	0	4 (16.7)
Thrombocytopenia	6 (8.5)	4 (16.7)	1 (4.3)	1 (4.2)
Nausea	4 (5.6)	1 (4.2)	3 (13.0)	0
Anemia	6 (8.5)	3 (12.5)	1 (4.3)	2 (8.3)
Lymphopenia	4 (5.6)	3 (12.5)	1 (4.3)	0
AST increased	4 (5.6)	3 (12.5)	1 (4.3)	0
ALT increased	2 (2.8)	2 (8.3)	0	0
Vomiting	3 (4.2)	1 (4.2)	2 (8.7)	0
Fatigue	4 (5.6)	1 (4.2)	2 (8.7)	1 (4.2)

All arms received 200mg abemaciclib monotherapy twice per day: Arm 1 = taken with a meal; Arm 2 = taken without a meal (modified fasting condition); Arm 3 = taken without regard to food. First three treatment cycles only. Abbreviations: ALT – alanine aminotransferase; AST – aspartate aminotransferase. N = number of patients receiving at least 1 abemaciclib dose (safety population); n = number of patients within category. All events occurring in \geq 2 patients in any one arm are included.

*One patient in treatment Arm 1 died due to respiratory failure and one patient in Arm 2 died due to abnormal hepatic function; neither death was considered related to the study treatment.

REFERENCES

1. Sledge, G.W., Jr., et al., *MONARCH 2: Abemaciclib in Combination With Fulvestrant in Women With HR+/HER2- Advanced Breast Cancer Who Had Progressed While Receiving Endocrine Therapy*. J Clin Oncol, 2017. **35**(25): p. 2875-2884.
2. FDA.gov. *Prescribing information VERZENIO™ (abemaciclib)*. 19th January 2020]; Available from: https://www.accessdata.fda.gov/drugsatfda_docs/label/2017/208716s000lbl.pdf.
3. Dickler, M.N., et al., *MONARCH 1, A Phase II Study of Abemaciclib, a CDK4 and CDK6 Inhibitor, as a Single Agent, in Patients with Refractory HR(+)/HER2(-) Metastatic Breast Cancer*. Clin Cancer Res, 2017. **23**(17): p. 5218-5224.
4. Sledge, G.W., Jr., et al., *The Effect of Abemaciclib Plus Fulvestrant on Overall Survival in Hormone Receptor-Positive, ERBB2-Negative Breast Cancer That Progressed on Endocrine Therapy- MONARCH 2: A Randomized Clinical Trial*. JAMA Oncol, 2020. **6**(1): p. 116-124.
5. Johnston, S., et al., *MONARCH 3 final PFS: a randomized study of abemaciclib as initial therapy for advanced breast cancer*. NPJ Breast Cancer, 2019. **5**: p. 5.
6. Goetz, M.P., et al., *Health-Related Quality of Life in MONARCH 3: Abemaciclib plus an Aromatase Inhibitor as Initial Therapy in HR+, HER2- Advanced Breast Cancer*. Oncologist, 2020. **25**(9): p. e1346-e1354.
7. Kaufman, P.A., et al., *Health-Related Quality of Life in MONARCH 2: Abemaciclib plus Fulvestrant in Hormone Receptor-Positive, HER2-Negative Advanced Breast Cancer After Endocrine Therapy*. Oncologist, 2020. **25**(2): p. e243-e251.
8. Turner, K., et al., *Abstract CT152: Food effect on the pharmacokinetics of 200-mg abemaciclib in healthy subject*. Cancer Research, 2016. **76**(14_Supplement): p. CT152-CT152.
9. Schmidt, L.E. and K. Dalhoff, *Food-drug interactions*. Drugs, 2002. **62**(10): p. 1481-502.
10. Chappell, J.C., et al., *Abemaciclib Inhibits Renal Tubular Secretion Without Changing Glomerular Filtration Rate*. Clin Pharmacol Ther, 2019. **105**(5): p. 1187-1195.
11. Schwartzberg, L., *Electronic Patient-Reported Outcomes: The Time Is Ripe for Integration Into Patient Care and Clinical Research*. American Society of Clinical Oncology Educational Book, 2016(36): p. e89-e96.
12. Rugo, H.S., et al., *The association of early toxicity and outcomes for patients treated with abemaciclib*. Journal of Clinical Oncology, 2018. **36**(15_suppl): p. 1053-1053.
13. Basch, E., et al., *Overall Survival Results of a Trial Assessing Patient-Reported Outcomes for Symptom Monitoring During Routine Cancer Treatment*. Jama, 2017. **318**(2): p. 197-198.
14. Basch, E., et al., *Symptom Monitoring With Patient-Reported Outcomes During Routine Cancer Treatment: A Randomized Controlled Trial*. J Clin Oncol, 2016. **34**(6): p. 557-65.

[Click here to view linked References](#)

1
2
3
4 **An open label, randomized phase 2 trial assessing the impact of food on the tolerability of**
5
6 **abemaciclib in patients with Advanced Breast Cancer**
7
8
9

10
11 **Authors:** Elgene Lim^{1#}, Frances Boyle^{2#}, Meena Okera³, Sherene Loi⁴, Sema Sezgin Goksu⁵, Gertjan
12 van Hal⁶, Sonya C. Chapman⁷, Jonathon Colby Gable⁸, Yanyun Chen⁸, Gregory L. Price⁸, Anwar M.
13 Hossain⁸, M. Corona Gainford⁸, Meritxell Bellet Ezquerro⁹
14
15
16
17

18 **Author Affiliations:** ¹Garvan Institute of Medical Research, St. Vincent's Clinical School, UNSW, NSW,
19 Australia; ²Mater Hospital, North Sydney, NSW, Australia; ³Adelaide Cancer Centre, Kurralta Park, SA,
20 Australia; ⁴Peter MacCallum Cancer Centre, Melbourne, VI, Australia; ⁵Akdeniz University Medical
21 Faculty, Antalya, Turkey; ⁶Eli Lilly and Company, Utrecht, The Netherlands; ⁷Eli Lilly and Company,
22 Windlesham, Surrey, UK; ⁸Eli Lilly and Company, Indianapolis, IN, USA; ⁹Hospital Universitario Vall
23 d'Hebron and Vall d'Hebron Institute of Oncology (VHIO), Barcelona, Spain.
24
25
26
27
28
29
30

31 # Contributed equally to the study
32
33

34 *** Corresponding Author:**
35

36 Elgene Lim
37
38

39 St. Vincent's Hospital Sydney,
40
41

42 Kinghorn Cancer Centre, 370 Victoria Street,
43
44

45 Darlinghurst, NSW, Australia 2010
46
47

48 e.lim@garvan.org.au
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 **ABSTRACT**
5
6

7 Purpose: Abemaciclib, a CDK4 & 6 inhibitor, is indicated for advanced breast cancer treatment. Diarrhea
8 is a frequently associated adverse event of abemaciclib. The study objective was to investigate if food
9 intake impacts local gastrointestinal toxicity.
10

11
12
13 Methods: This Phase 2 study (I3Y-MC-JPCP, NCT03703466) randomized 72 patients 1:1:1 to receive
14 abemaciclib 200mg monotherapy twice daily (1) with a meal, (2) in a modified fasting state or (3) without
15 regard to food. Primary endpoints included: incidence of investigator assessed severe (\geq Grade 3),
16 prolonged (>7 -days) Grade 2 diarrhea, treatment discontinuation, dose modifications, and loperamide
17 utilization during the first 3 cycles of treatment. Patient outcomes were captured via a daily electronic
18 diary. Pharmacokinetics (PK) are reported.
19
20
21
22
23
24
25

26 Results: Incidence of investigator assessed severe diarrhea (Grade ≥ 3) was 1.4% (1 patient in Arm 1).
27 Median duration of Grade 3 diarrhea was 1 day by both investigator assessment (1 patient in Arm 1) and
28 patient-reported assessment (1 patient each in Arms 1 and 3). Median duration of investigator-assessed
29 Grade 2 diarrhea was 2 days overall. No patient discontinued treatment due to diarrhea. Nine patients
30 (12.7%) had a dose reduction, and 7 patients (9.9%) had a dose omission due to diarrhea. Ninety-four
31 percent of patients used loperamide at least once. Abemaciclib PK was comparable across the 3 arms.
32
33
34
35
36
37
38

39 Conclusion: The results suggest that diarrhea incidence associated with abemaciclib was unrelated to
40 timing of food intake, was predominantly low grade, of short duration and well managed with loperamide
41 and dose modifications.
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

INTRODUCTION

Abemaciclib is an oral, selective, and potent inhibitor of cyclin-dependent kinases 4 and 6 (CDK4 & 6) dosed twice daily (BID) on a continuous schedule [1]. Resulting from the MONARCH series of clinical trials, abemaciclib is approved as monotherapy and in combination with endocrine therapy (ET) for the treatment of patients with hormone receptor-positive (HR+), human epidermal growth factor receptor 2-negative (HER2-) advanced breast cancer (ABC) [2].

MONARCH 1, a single-arm Phase 2 study of abemaciclib 200mg BID monotherapy in patients with refractory HR+, HER2- ABC, demonstrated promising clinical activity (objective response rate (ORR) of 19.7% (95% confidence interval [CI]: 13.3-27.5) [3].

MONARCH 2 was a randomized, double-blind, Phase 3 study of abemaciclib 150mg BID in combination with fulvestrant compared to placebo plus fulvestrant in women with HR+, HER2- ABC who had progressed following ET therapy [1]. This trial demonstrated significantly improved progression-free survival (PFS) (median 16.4 versus 9.3 months; hazard ratio [HR] [95% CI]: 0.553 [0.449-0.681]; $p < .001$) and overall survival (OS) (median 46.7 versus 37.3 months; HR [95% CI]: 0.757 [0.606-0.945]; $p = .01$) [1, 4].

MONARCH 3 was a randomized, double-blind Phase 3 study of abemaciclib 150mg BID in combination with a non-steroidal aromatase inhibitor (NSAI) compared to placebo plus NSAI as initial therapy in women with HR+, HER2- ABC [5]. Abemaciclib plus NSAI significantly improved PFS (median 28.18 versus 14.76 months; HR [95% CI]: 0.540 [0.418-0.698]; $p = .000002$).

Diarrhea was the most frequently reported treatment-emergent adverse event (TEAE) of any-grade in patients prescribed abemaciclib in all three pivotal studies, irrespective of whether abemaciclib was taken as a monotherapy (MONARCH 1 [200mg BID]: diarrhea=90.2%) or in combination with ET (MONARCH 2 and MONARCH 3 [150mg BID]: diarrhea=87.1% and 82.3% respectively) [1, 3, 5]. Grade 3 diarrhea was reported in 20% of patients in MONARCH 1, 14% in MONARCH 2 and 10% in MONARCH 3 [3-5]. In all three trials, incidence of Grade 2 and Grade 3 diarrhea was greatest during the first month of treatment and decreased over the remaining cycles of therapy (Fig. 1). The median duration of any-grade diarrhea

1
2
3
4 was similar across the 3 trials ranging 6-8 days, with the median duration of Grade 2 diarrhea ranging 8-
5
6 11 days and Grade 3 diarrhea, 5-8 days [1, 3, 5]. In each study, diarrhea was retrospectively assessed by
7
8 the investigator at the beginning of each 28-day cycle and graded as per Common Terminology Criteria
9
10 for Adverse events (CTCAE) criteria [1, 3, 5]. To our knowledge, no patient-reported daily data on
11
12 abemaciclib and diarrhea have been published to date.

13
14
15 Standardized anti-diarrheal management plans are outlined in the product label [2]. Patients are advised
16
17 to commence over-the-counter (OTC) antidiarrheal medication such as loperamide at the first sign of
18
19 loose stools, increase their fluid intake, and contact their physician. If diarrhea has not resolved within 24
20
21 hours, abemaciclib should be suspended until resolution occurs. Resolution is defined as either a
22
23 reduction to baseline or Grade 1 (<4 stools per day increase over baseline). The label also includes
24
25 detailed guidance for dose modifications and reductions according to severity of diarrhea [2].
26
27

28 Consistent with management guidance, 21%, 19%, and 14% of patients in MONARCH 1, 2, and 3
29
30 respectively required dose reductions and 1%, 3%, and 2% respectively discontinued the study drug due
31
32 to diarrhea [1, 3, 5]. Antidiarrheal use across the studies varied between 61% in MONARCH 1, 76% in
33
34 MONARCH 2, and 61% in MONARCH 3.
35

36
37 In MONARCH 2 and 3, a paper version of the patient-reported outcomes (PRO) assessment was
38
39 completed once at baseline, and on treatment; with more frequent collection in earlier cycles, and at the
40
41 follow-up visit [6, 7]. PRO results from abemaciclib in combination with fulvestrant (MONARCH 2) or in
42
43 combination with NSAID (MONARCH 3) did not show clinically significant differences in patient-reported
44
45 global health, functioning, or most symptoms compared to ET alone [6, 7]. Baseline scores were similar
46
47 between treatment arms in each study. In both trials, diarrhea was the only patient-reported symptom with
48
49 a statistically significant and clinically meaningful difference between treatment arms. These diarrhea
50
51 findings were reported in early treatment cycles, consistent with investigator assessments, decreased in
52
53 later cycles and returned to near baseline levels at the post-therapy follow-up visits [6, 7].
54

55
56 For some drugs, coadministration with food can impact bioavailability and may have clinically significant
57
58 consequences. In clinical studies, a high-fat, high-calorie meal increased the exposure (AUC) of
59
60 abemaciclib analytes by 9% and increased C_{max} by 25% [8]. These changes in exposure are not clinically
61

1
2
3
4 meaningful and abemaciclib is therefore given without regard to food. However, it is possible that taking
5
6 abemaciclib with food may impact local gastrointestinal toxicity independently of systemic
7
8 pharmacokinetics (PK) and thus alter drug tolerability. As an example, ingestion of food with nonsteroidal
9
10 anti-inflammatory drugs (NSAIDs) is often preferred because it reduces local gastrointestinal adverse
11
12 effects [9]. In order to address this issue and at the request of the U.S. Food and Drug Administration
13
14 (FDA), this randomized study (I3Y-MC-JPCP) evaluated the impact of coadministration of food on the
15
16 incidence and tolerability of diarrhea in patients with HR+, HER2- ABC receiving abemaciclib
17
18 monotherapy.
19
20
21
22

23 **METHODS**

24 *Study design and patients*

25
26
27 JPCP (NCT03703466) was a global, randomized, open-label Phase 2 study evaluating the timing of food
28
29 intake on the incidence of severe diarrhea (Grade ≥ 3) or prolonged Grade 2 diarrhea (>7 days duration)
30
31 when receiving abemaciclib monotherapy 200mg orally (PO) BID in patients with previously treated HR+,
32
33 HER2- ABC.
34
35
36
37

38 The study was conducted at 15 centers in 5 countries (Australia, Belgium, Russian Federation, Spain and
39
40 Turkey). It was approved by ethical and local institutional review boards for the participating sites and was
41
42 conducted according to the Declaration of Helsinki. Patients provided written informed consent prior to
43
44 trial enrollment. The study was overseen by an ethics review board.
45
46

47 Eligible patients included males and females ≥ 18 years of age with a diagnosis of recurrent, locally
48
49 advanced, unresectable, or metastatic HR+, HER2- ABC. Patients were required to have an Eastern
50
51 Cooperative Oncology Group (ECOG) performance status of ≤ 1 ; a willingness to use an e-diary; and
52
53 having no prior use of a CKD4 & 6 inhibitor. Patients must have received ≥ 1 chemotherapy regimen and
54
55 progressed after prior anti-estrogen therapy for ABC. Patients were required to have discontinued all
56
57 previous treatments for cancer and recovered from the acute effects of therapy. Per protocol, length of
58
59 time between end of previous treatment and first abemaciclib dose was 14 to 28 days, depending on type
60
61
62
63
64
65

1
2
3
4 of prior treatment. Patients were ineligible if they had a serious concomitant systemic disorder (for
5 example, active infection or a gastrointestinal disorder causing clinically significant symptoms such as
6 nausea, vomiting or diarrhea [such as Crohn's disease, ulcerative colitis], or profound immune
7 suppression) or a serious preexisting medical condition (for example, history of major surgical resection
8 involving the stomach or small bowel) that, in the opinion of the investigator, would compromise/preclude
9 the patient's ability to adhere to the protocol.
10
11
12
13
14
15

16
17 Patients were randomly assigned 1:1:1 to take abemaciclib 200mg monotherapy either; with a meal (Arm
18 1); in a modified fasting state defined as at least 1 hour before or 2 hours after a meal (Arm 2); or without
19 regard to food (Arm 3). A meal was defined as whatever the patient would normally eat at that time.
20
21
22

23 Patients were advised to avoid the consumption of grapefruit or grapefruit juice and other inducers and
24 inhibitors of cytochrome P450 (CYP) 3A where possible, as these can affect the exposure of abemaciclib.
25
26

27 Abemaciclib was administered on a continuous BID schedule with at least 6 hours separating doses.
28

29 Treatment cycles lasted 28 days. The study period consisted of the first 3 cycles of treatment for each
30 individual patient and all results presented herein reflect the first 3 cycles only. Patients who continued to
31 receive benefit following cycle 3 remained on treatment at investigator discretion and took abemaciclib
32 without regard to food as per label. All study procedures were followed until study completion, which
33 occurred when the last enrolled patient completed 3 cycles. Loperamide was the protocol specified anti-
34 diarrheal medication. Patients received diarrhea management guidance per label and were provided with
35 written support materials.
36
37
38
39
40
41
42
43

44 A training program using both face-to-face guidance and virtual media was developed to ensure
45 standardized implementation of the e-diary globally. All patients completed a daily e-diary detailing the
46 timing of each abemaciclib dose in relation to a meal, number of bowel movements (BM), and number of
47 loperamide tablets. Supplemental Fig. 1 visualizes the patient's e-diary experience and how they were
48 prompted to respond within the e-diary. To determine an accurate baseline assessment, patients
49 recorded daily number of BM for a week prior to study commencement. Adherence to study medication
50 and e-diary was made available for investigator and site staff review.
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 PK samples were collected prior to first dose on cycle 1 day 1 and then in conjunction with other
5
6 laboratory samples in cycle 1: day 15, cycle 2: days 1 and 15, and cycle 3: day 1.
7

8 9 *Objectives*

10
11 The primary objective was to evaluate and summarize investigator-assessed incidence of Grade ≥ 3 and
12 prolonged Grade 2 (>7 days continuous duration) diarrhea; dose reductions, interruptions, and
13 discontinuations due to diarrhea; and patient-reported utilization of anti-diarrheal medications during the
14 first 3 cycles of treatment. Secondary objectives included overall safety, incidence and severity of TEAEs,
15 serious adverse events, deaths and clinical laboratory abnormalities. PK analysis included steady-state
16 concentrations of abemaciclib, its active metabolites LSN2839567 (M2) and LSN3106726 (M20), and
17 total active analytes (sum of abemaciclib+M2+M20). An exploratory objective was to evaluate and
18 summarize incidence and duration of diarrhea reported daily by the patient using an e-diary.
19
20
21
22
23
24
25
26
27

28 29 *Statistical Analysis*

30
31 The study planned to enroll approximately 60 patients and was descriptive in nature. It was not powered
32 for formal statistical comparison between groups, and sample size was based on regulatory guidance.
33
34 Assignment to treatment arms were determined by a computer-generated random sequence and
35 randomization was stratified by region.
36
37
38
39

40 Baseline analyses and patient dispositions were based on the intent-to-treat (ITT) population which
41 included all patients enrolled and randomized in the trial. Safety analyses included all patients who
42 received at least 1 dose of abemaciclib. Analyses of all primary investigator-assessed endpoints and
43 adverse events were based on the safety population and summarized by study arms. Investigator
44 assessments of diarrhea and dose modifications were required on day 1 of each cycle and were
45 conducted as per their usual practice. While investigators and site staff had real time access to e-diary
46 data to facilitate monitoring, ensure timely patient education, and for the grading of diarrhea, investigators
47 were not required to review the e-diary data in their assessment of diarrhea. Adherence with study
48 medication and e-diary was monitored and recorded by investigators. Patient data collected via the e-
49 diary was summarized descriptively to provide daily estimates of loperamide use, compliance with
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 assignment to food administration cohort, as well as incidence and duration of diarrhea, complementary
5
6 to investigator assessment. PK analysis included all patients who had received at least 1 dose of
7
8 abemaciclib and who had at least 1 evaluable PK sample.
9

10 11 12 13 **RESULTS**

14 15 16 *Patients*

17
18
19 Seventy-two patients were enrolled from December 4th, 2018 to April 10th, 2019 with 24 patients
20
21 randomized to each of the 3 arms (Fig. 2). One patient in Arm 2 withdrew from the study and did not
22
23 receive treatment, resulting in a safety population of 71 patients. Only 1 patient was male; this patient was
24
25 randomized to Arm 1 (Table 1). The majority of patients (71%) were recruited from Australia, Spain, and
26
27 Belgium. The mean age (years) was similar in Arm 1 (58), Arm 2 (57), and Arm 3 (58). Approximately
28
29 22% of patients had received ≥ 5 prior chemotherapies for ABC. Mean treatment compliance calculated by
30
31 pill return across the 3 groups was 95.1%. Mean compliance with the assigned arm's timing of
32
33 abemaciclib dosing in relation to a meal was 97.4%. Mean e-diary completion compliance across all
34
35 study arms was 95.7%.
36

37
38 Although only 2 of 18 investigators assessed the e-diary portal 3 or more times during the study period,
39
40 the portal was accessed more frequently by study coordinators and nurses at each site. At the July 8th,
41
42 2019 data cutoff, 30 patients (41.7%) had discontinued treatment, with 21 patients (29.2%) discontinuing
43
44 due to progressive disease.
45

46 47 *Diarrhea incidence and dose modifications*

48
49
50 Food did not appear to impact incidence of investigator assessed diarrhea or dose modifications. While
51
52 the numbers were low, the incidence of Grade ≥ 3 and prolonged Grade 2 diarrhea during the first 3 cycles
53
54 of treatment were comparable across the 3 arms (Table 2). Notably, investigator assessed Grade 3
55
56 diarrhea occurred in only 1 patient overall (1.4%) during cycle 1 in Arm 1 and lasted 1 day. Prolonged
57
58 Grade 2 diarrhea (>7 -days) occurred in 11 patients overall (15.5%).
59
60
61
62
63
64
65

1
2
3
4 Dose reductions and omissions (dose interruptions) were balanced across Arms 1, 2, and 3 with 9
5
6 (12.7%) of all 71 patients having at least 1 dose reduction due to diarrhea and 7 (9.9%) of all patients
7
8 requiring a dose omission for diarrhea (Table 2). 94% of patients used loperamide at least once in the first
9
10 3 cycles. Dose reduction due to fatigue and neutropenia were also balanced across all 3 arms, occurring
11
12 in 4 (5.6%) and 8 (11.3%) of all 71 patients respectively.
13

14 *Pharmacokinetics*

15
16 A total of 69 patients had evaluable PK plasma samples: 24, 21, and 24 in Arms 1, 2, and 3 respectively.
17
18 A total of 260, 256, and 259 samples were available for abemaciclib, and its active metabolites M2 and
19
20 M20 respectively. Abemaciclib concentrations were comparable across the 3 treatment arms (Fig. 3), with
21
22 average values ranging from 305 to 369 ng/ml on cycle 1 day 15. The mean range for total active
23
24 analytes for the same time point was similarly comparable (1.38–1.60 uM).
25
26

27 *Treatment-Emergent Adverse Events (TEAEs)*

28
29 Investigator-reported TEAEs of diarrhea were experienced by 21 patients (87.5%) in Arm 1, 22 patients
30
31 (95.7%) in Arm 2, and 22 patients (91.7%) in Arm 3. No Grade 4 or 5 diarrhea events were reported
32
33 (Table 2). TEAEs were otherwise consistent with prior abemaciclib trials and are listed in Table 3. No
34
35 patients experienced a serious AE of diarrhea, or discontinued abemaciclib due to diarrhea, during the
36
37 first 3 cycles. Importantly, despite the frequent use of loperamide, no patient experienced Grade 3 or
38
39 greater constipation. Overall, 2 patients experienced Grade 1 constipation and 1 patient had a Grade 2
40
41 event. Two deaths occurred while on therapy or within 30 days of treatment discontinuation, both of which
42
43 were associated with the study disease and assessed by the investigators to be unrelated to the study
44
45 treatment. One death (4.2%) in Arm 1 was due to respiratory failure related to pulmonary progressive
46
47 disease and pulmonary infection, and 1 death (4.3%) in Arm 2 was due to abnormal hepatic function
48
49 where the patient had been diagnosed with liver metastasis prior to starting treatment. The most common
50
51 treatment-emergent laboratory toxicities experienced by all patients included increased creatinine and
52
53 decreased neutrophil count. Increases in serum creatinine levels are a known pharmacodynamic effect of
54
55 abemaciclib caused by inhibition of renal tubular secretion without affecting glomerular function [10]. The
56
57
58
59
60
61
62
63
64
65

1
2
3
4 TEAE of neutropenia has been observed across MONARCH studies. Neutropenia was not associated
5
6 with severe infection [1, 3, 5].
7
8

9 Diarrhea incidence and duration as assessed by the investigator (A and B) and the patient (C and D)
10 during the first 3 cycles of therapy is illustrated in Fig. 4. This visually highlights the discrepancy between
11 both assessments. One episode of Grade 3 diarrhea was reported by investigator assessment (for 1
12 patient in Arm 1) while 2 episodes were reported by patient assessment (1 patient each in Arm 1 and Arm
13 3). All reported Grade 3 episodes were of 1 day's duration. Grade 2 diarrhea as assessed by the
14
15 investigator was of longer duration than that assessed by the patient (median duration 2 days vs 1 day).
16
17 Median duration of investigator-assessed Grade 2 diarrhea was 2 days overall: 2 days (Arm 1), 7 days
18 (Arm 2), and 3 days (Arm 3). Finally, Fig. 4A demonstrates that Grade 1 diarrhea was assessed by the
19 investigator as continuous while patient daily assessment via the e-diary (Fig. 4C) reported intermittent
20 and short duration Grade 1.
21
22
23
24
25
26
27
28

29 Loperamide doses and corresponding patient-reported grade of diarrhea (calculated from daily numbers
30 of BM relative to baseline recorded in the patient e-diaries) are displayed in Fig. 4E. While loperamide
31 was used frequently, the number of tablets taken varied daily with the grade of diarrhea experienced.
32
33
34
35
36
37
38

39 **DISCUSSION**

40
41
42 Results from this trial demonstrate that food does not appear to impact the incidence or duration of
43 prolonged Grade 2 and Grade ≥ 3 diarrhea, supporting the product label that abemaciclib may be taken
44 with or without food. Abemaciclib PK was comparable across the 3 treatment arms, indicating no major
45 differences in drug exposure related to meal timing. It has been reported that a high-fat, high-calorie meal
46 increases abemaciclib plasma concentrations to a non-clinically meaningful extent [8]. However, the
47 conditions of this trial simulate real-world drug administration rather than the high-fat, high-calorie meals
48 in clinical studies specifically designed to assess the effect of food on PK parameters.
49
50
51
52
53
54
55

56
57 The findings reported here include several novel insights about abemaciclib-associated diarrhea beyond
58 evaluation of the effect of food. Electronic capture of patient-reported outcomes (ePRO) can provide more
59
60
61

1
2
3
4 granular reporting of adverse events than can be captured by traditional techniques [11]. Investigator
5 assessments are complemented by daily patient reports of BM frequency and antidiarrheal use recorded
6 with easy-to-use handheld e-diaries (Figs 4A and B). Capturing daily patient-reported data via e-diary
7 facilitated detailed, real-time characterization of the patient experience compared to retrospective
8 physician assessment alone. Mean compliance with daily completion of the e-diary was 96% and is
9 remarkable for a global trial in a patient population with heavily pre-treated ABC. This suggests that it is
10 feasible, with appropriate training and oversight, to employ such tools in oncology trials. The overall
11 incidence of diarrhea (any-grade) in the current study was similar to that in MONARCH 1, where the dose
12 used, and patient population were the same. However, the incidence of Grade 3 diarrhea was much lower
13 here (1% vs 20%). This could suggest that as physicians have gained more experience with abemaciclib
14 and incorporated standard diarrhea management guidelines into clinical care, abemaciclib-associated
15 diarrhea is better understood and better managed. An assessment of the baseline number of BM as
16 performed in this study would be important to allow an accurate grading of diarrhea as per CTCAE
17 criteria. The median duration of Grade 2 and Grade 3 diarrhea, reported by both investigators and
18 patients, was shorter than what was observed in the MONARCH 1, 2, and 3 trials [3-5]. Interestingly,
19 patients in this study reported earlier resolution of Grade 2 diarrhea than investigators (Fig. 4).
20 Furthermore, Fig. 4 illustrates that patients reported intermittent and shorter durations of Grade 1 diarrhea
21 as opposed to the continuous Grade 1 diarrhea assessed by investigators. These data demonstrate the
22 benefits of accurately recording grade and duration of diarrhea and reinforces the importance of recording
23 the patient experience in real time. The additional granularity obtained through use of the daily e-diary
24 permitted exploration of the relationship between antidiarrheal use and the incidence, grade, and duration
25 of diarrhea. While loperamide was frequently used over the course of the study (Fig. 4C), the pattern of
26 use suggests patients were appropriately educated regarding diarrhea management and varied their use
27 of loperamide according to the severity experienced.

28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54 Consistent with management guidelines in the product label allowing for dose modifications for diarrhea,
55 12.7% of patients in this study had at least 1 dose reduction. A previous exploratory analysis
56 demonstrated no difference in PFS outcomes for patients who dose reduced versus those who did not in
57 the MONARCH 1, 2, and 3 trials, suggesting patients can receive a reduced dose and still derive
58
59
60
61

1
2
3
4 meaningful benefit from abemaciclib treatment [12]. Those findings and the results of this study
5
6 collectively suggest the standardized diarrhea management plan described in the product label
7
8 appropriately manages abemaciclib-associated diarrhea without risking a decrease in efficacy.
9

10
11 This study had several limitations. First, it was of modest size limiting the ability to make comparisons
12
13 between arms. Additionally, while data collected via e-diary was made available to site staff in real-time,
14
15 e-diary user metrics reports showed it was predominantly accessed by study coordinators and nurses
16
17 with only 2 of 18 investigators accessing the data ≥ 3 times in the 3-month study period (data on file). Ten
18
19 of 18 investigators never accessed the e-diary data. This potentially explains the discrepancies seen
20
21 between investigator and patient assessments (Fig. 4). Although high compliance rates with the daily
22
23 patient e-diaries provided valuable insights into diarrhea incidence and management, it could be argued
24
25 that patients may have been biased towards better management because of increased awareness and a
26
27 daily requirement to enter data. Prior studies have indicated that active assessment and monitoring of
28
29 symptomatic adverse events could yield positive outcomes for patients, including health related quality of
30
31 life [13, 14]. Further studies are needed to understand the impact of patient self-monitoring on outcomes.
32
33 Loperamide use was higher than what was previously seen in the MONARCH studies and it is possible
34
35 that the high use of anti-diarrheal medication masked the impact of food on the incidence and severity of
36
37 diarrhea. Food composition intake was also not standardized across groups. Finally, while PK analysis
38
39 detected no discernable differences between the 3 arms, the study was not designed to assess the
40
41 impact of food on abemaciclib PK, and it is likely that only large effects would be detected with a non-
42
43 crossover study design and sparse PK sampling.
44

45
46 In conclusion, the results suggest that abemaciclib can be taken with or without food. Management
47
48 guidance for associated diarrhea provided in the product label is appropriate. The current study results
49
50 suggest that diarrhea is manageable with loperamide and dose modifications as needed. This is
51
52 demonstrated by a progressive decrease in the frequency of Grade 3 diarrhea (in the first 3 cycles) for
53
54 MONARCH 1, 2, and 3, and in the present study. Diarrhea onset is usually in the first cycle, and patient-
55
56 reported e-diary data in the present study shows shorter duration of Grade 1 and 2 diarrhea than reported
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

by investigator assessment. It is critical that patients receiving abemaciclib are educated on how to manage diarrhea and commence loperamide at the first sign of loose stools for optimal treatment benefit.

1
2
3
4 **DATA AVAILABILITY**
5

6
7 Eli Lilly and Company makes patient-level data available from Lilly-sponsored studies on marketed drugs
8
9 for approved uses following acceptance for publication. Lilly is one of several companies that provide this
10
11 access through the website clinicalstudydatarequest.com. Qualified researchers can submit research
12
13 proposals and request anonymized data to test new hypotheses. Lilly's data sharing policies are provided
14
15 on the clinicalstudydatarequest.com site under the Study Sponsors page.
16
17

18
19
20 **ACKNOWLEDGEMENTS**
21

22 The authors are grateful to the patients, their families, and caregivers for participating in study JPCP, and
23
24 to the study investigator and site staff for their collaboration. Writing and editing provided by Nicholas
25
26 Pulliam, Sandra Deady and John Hurley of Eli Lilly and Company. Funding provided by Eli Lilly and
27
28 Company.
29
30

31
32 **AUTHOR CONTRIBUTIONS**
33

34 All authors contributed to the intellectual content of the work through conception, design, data acquisition,
35
36 analysis and interpretation as well as the drafting and critical review of the manuscript. All authors
37
38 provided their final approval of the manuscript.
39
40
41

42 **COMPETING INTERESTS INFORMATION**
43

44
45 EL reports a consulting or advisory (paid to his institution) role with Eli Lilly and Company, Novartis
46
47 Australia, Pfizer and Roche; institutional research funding from Novartis Australia; travel or
48
49 accommodation expenses from Eli Lilly and Company; and a patent/royalty/intellectual property for the
50
51 sensitization of BCL-2-expressing breast tumors to chemotherapy by the BH3 mimetic ABT-737. FB
52
53 reports a consulting or advisory role with Pfizer, Eli Lilly and Company, Novartis Australia, Roche and
54
55 Eisai; and travel or accommodation expenses from Novartis Australia. MO has no competing interests to
56
57 declare. SL reports receiving research funding to her institution from Novartis, Bristol Meyers Squibb,
58
59 Merck, Puma Biotechnology, Eli Lilly, Nektar Therapeutics Astra Zeneca, Roche-Genentech and Seattle
60
61
62
63
64
65

1
2
3
4 Genetics; a consulting or advisory role (not compensated) with Seattle Genetics, Novartis, Bristol Meyers
5
6 Squibb, Merck, AstraZeneca and Roche-Genentech; a consulting or advisory role (paid to her institution)
7
8 with Aduro Biotech, Novartis, GlaxoSmithKline, Roche-Genentech, Astra Zeneca, Silverback
9
10 Therapeutics, G1 Therapeutics, PUMA Biotechnologies, Pfizer, Seattle Genetics and Bristol Meyers
11
12 Squibb; is a Scientific Advisory Board Member of Akamara Therapeutics; and is supported by the
13
14 National Breast Cancer Foundation of Australia Endowed Chair and the Breast Cancer Research
15
16 Foundation, New York. SSG reports a consulting or advisory role with Novartis, MSD, Roche and Amgen;
17
18 speakers' bureau company with Eli Lilly and Company, Novartis, Roche, Phizer and BMS; travel,
19
20 accomodation or congress fees from Pfizer, Novartis, Roche, Astra-Zeneca; and is a primary/sub
21
22 investigator of many clinical studies sponsored by Roche, MSD, BMS and Novartis. SCC, JCG, YC, GLP,
23
24 AMH and MCG are full-time employees of Eli Lilly and Company and are Eli Lilly and Company
25
26 shareholders. MB-E reports personal fees from Pfizer, Novartis, and Eli Lilly and Company; and travel
27
28 expenses from Pfizer and Roche.
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figure Legends

Fig. 1 Comparison of investigator-assessed treatment-emergent diarrhea in patients receiving abemaciclib from studies JPCP, MONARCH 1, MONARCH 2, and MONARCH 3

Fig. 2 Consort diagram

Fig. 3 Abemaciclib concentrations on Cycle 1 Day 15 across the three treatment arms

Fig. 4 Duration and incidence of diarrhea in cycles 1-3

- A. Investigator assessed
- B. Investigator assessed
- C. Patient-reported e-diary
- D. Patient-reported e-diary
- E. Patient-reported e-diary account of loperamide doses in cycles 1-3

Suppl Fig. 1 Patient facing e-diary

Table Legends

Table 1 Demographics and baseline characteristics

Table 2 Primary endpoint results, investigator assessed

Table 3 Treatment-emergent adverse events (TEAEs) \geq Grade 3; investigator-assessed

Table 1 Demographics and baseline characteristics

	Overall	Arm 1	Arm 2	Arm 3
	N = 72	N = 24	N = 24	N = 24
Sex, n (%)				
Female	71 (98.6)	23 (95.8)	24 (100.0)	24 (100.0)
Male	1 (1.4)	1 (4.2)	0	0
Age categories, n (%)				
<65 years	53 (73.6)	17 (70.8)	15 (62.5)	21 (87.5)
≥65 years	19 (26.4)	7 (29.2)	9 (37.5)	3 (12.5)
Age (years)				
Mean	57.6	58.2	56.9	57.6
Race, n (%)				
Asian	2 (2.8)	2 (8.3)	0	0
White	70 (97.2)	22 (91.7)	24 (100.0)	24 (100.0)
BMI, kg/m² (SD)				
Mean	26.9 (4.5)	26.6 (4.3)	28.3 (5.4)	26.0 (3.6)
Region, n (%)				
Turkey / Russia	21 (29.2)	7 (29.2)	7 (29.2)	7 (29.2)
Australia / Spain / Belgium	51 (70.8)	17 (70.8)	17 (70.8)	17 (70.8)
ECOG Performance Status, n (%)				
0	37 (51.4)	13 (54.2)	10 (41.7)	14 (58.3)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1	34 (47.2)	11 (45.8)	13 (54.2)	10 (41.7)
Nature of disease, n (%)				
Visceral	59 (81.9)	20 (83.3)	17 (70.8)	22 (91.7)
Bone only	8 (11.1)	3 (12.5)	5 (20.8)	0
Others	4 (5.6)	1 (4.2)	1 (4.2)	2 (8.3)
Missing	1 (1.4)	0	1 (4.2)	0
Number of organ sites (%)				
1	14 (19.4)	3 (12.5)	8 (33.3)	3 (12.5)
2	24 (33.3)	9 (37.5)	5 (20.8)	10 (41.7)
≥3	33 (45.8)	12 (50.0)	10 (41.7)	11 (45.8)
Number of prior chemotherapy regimens for ABC, n (%)				
<5	54 (75.0)	20 (83.3)	14 (60.9)	20 (83.3)
≥5	16 (22.2)	4 (16.7)	9 (39.1)	3 (12.5)

Intention-to-treat (ITT) population. All arms received 200mg abemaciclib monotherapy twice per day: Arm 1 = taken with a meal; Arm 2 = taken without a meal (modified fasting condition); Arm 3 = taken without regard to food. Abbreviations: ECOG = Eastern Cooperative Oncology Group; N = number of patients in the intent-to-treat population; n = number of patients within category. ECOG status missing for 1 patient in Arm 2

Table 2 Primary endpoint results, investigator assessed

	Overall	Arm 1	Arm 2	Arm 3
	N = 71	N = 24	N = 23*	N = 24
Endpoint				
Diarrhea (any grade), n (%)	65 (91.5)	21 (87.5)	22 (95.7)	22 (91.7)
Grade 1 diarrhea, n (%)	35 (49.3)	10 (41.7)	14 (60.9)	11 (45.8)
Duration of Grade 1 diarrhea, median days	8	9	6	10
Grade 2 diarrhea, n (%)	29 (40.8)	10 (41.7)	8 (34.8)	11 (45.8)
Duration of Grade 2 diarrhea, median days	2	2	7	3
Grade 2 diarrhea lasting >7 days, n (%)	11 (15.5)	2 (8.3)	4 (17.4)	5 (20.8)
Grade 3 diarrhea, n (%)	1 (1.4)	1 (4.2)	0	0
Duration of Grade 3 diarrhea, median days	1	1	0	0
Grade 4 diarrhea, n (%)	0	0	0	0
≥1 Dose reduction due to diarrhea, n (%)	9 (12.7)	4 (16.7)	2 (8.7)	3 (12.5)
≥1 Dose omission due to diarrhea, n (%)	7 (9.9)	4 (16.7)	1 (4.3)	2 (8.3)
Treatment discontinued due to diarrhea, n (%)	0	0	0	0

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Loperamide use, n (%)	67 (94.3)	23 (95.8)	21 (91.3)	23 (95.8)
-----------------------	-----------	-----------	-----------	-----------

All arms received 200mg abemaciclib monotherapy twice per day: Arm 1 = taken with a meal; Arm 2 = taken without a meal (modified fasting condition); Arm 3 = taken without regard to food. First three treatment cycles only. N = number of patients receiving at least 1 abemaciclib dose (safety population); n = number of patients within category

*one patient in Arm 2 discontinued from the study prior to receiving treatment

Table 3 Treatment-emergent adverse events (TEAEs) Grade ≥ 3 ; investigator-assessed

	Overall	Arm 1	Arm 2	Arm 3
	N = 71	N = 24	N = 23	N = 24
TEAE	n (%)	n (%)	n (%)	n (%)
Neutropenia	20 (28.2)	7 (29.2)	3 (13.0)	10 (41.7)
Leukopenia	9 (12.7)	5 (20.8)	0	4 (16.7)
Thrombocytopenia	6 (8.5)	4 (16.7)	1 (4.3)	1 (4.2)
Nausea	4 (5.6)	1 (4.2)	3 (13.0)	0
Anemia	6 (8.5)	3 (12.5)	1 (4.3)	2 (8.3)
Lymphopenia	4 (5.6)	3 (12.5)	1 (4.3)	0
AST increased	4 (5.6)	3 (12.5)	1 (4.3)	0
ALT increased	2 (2.8)	2 (8.3)	0	0
Vomiting	3 (4.2)	1 (4.2)	2 (8.7)	0
Fatigue	4 (5.6)	1 (4.2)	2 (8.7)	1 (4.2)


All arms received 200mg abemaciclib monotherapy twice per day: Arm 1 = taken with a meal; Arm 2 = taken without a meal (modified fasting condition); Arm 3 = taken without regard to food. First three treatment cycles only. Abbreviations: ALT – alanine aminotransferase; AST – aspartate aminotransferase. N = number of patients receiving at least 1 abemaciclib dose (safety population); n = number of patients within category. All events occurring in ≥ 2 patients in any one arm are included.

*One patient in treatment Arm 1 died due to respiratory failure and one patient in Arm 2 died due to abnormal hepatic function; neither death was considered related to the study treatment.


REFERENCES

1. Sledge, G.W., Jr., et al., *MONARCH 2: Abemaciclib in Combination With Fulvestrant in Women With HR+/HER2- Advanced Breast Cancer Who Had Progressed While Receiving Endocrine Therapy*. *J Clin Oncol*, 2017. **35**(25): p. 2875-2884.
2. FDA.gov. *Prescribing information VERZENIO™ (abemaciclib)*. 19th January 2020]; Available from: https://www.accessdata.fda.gov/drugsatfda_docs/label/2017/208716s000lbl.pdf.
3. Dickler, M.N., et al., *MONARCH 1, A Phase II Study of Abemaciclib, a CDK4 and CDK6 Inhibitor, as a Single Agent, in Patients with Refractory HR(+)/HER2(-) Metastatic Breast Cancer*. *Clin Cancer Res*, 2017. **23**(17): p. 5218-5224.
4. Sledge, G.W., Jr., et al., *The Effect of Abemaciclib Plus Fulvestrant on Overall Survival in Hormone Receptor-Positive, ERBB2-Negative Breast Cancer That Progressed on Endocrine Therapy- MONARCH 2: A Randomized Clinical Trial*. *JAMA Oncol*, 2020. **6**(1): p. 116-124.
5. Johnston, S., et al., *MONARCH 3 final PFS: a randomized study of abemaciclib as initial therapy for advanced breast cancer*. *NPJ Breast Cancer*, 2019. **5**: p. 5.
6. Goetz, M.P., et al., *Health-Related Quality of Life in MONARCH 3: Abemaciclib plus an Aromatase Inhibitor as Initial Therapy in HR+, HER2- Advanced Breast Cancer*. *Oncologist*, 2020. **25**(9): p. e1346-e1354.
7. Kaufman, P.A., et al., *Health-Related Quality of Life in MONARCH 2: Abemaciclib plus Fulvestrant in Hormone Receptor-Positive, HER2-Negative Advanced Breast Cancer After Endocrine Therapy*. *Oncologist*, 2020. **25**(2): p. e243-e251.
8. Turner, K., et al., *Abstract CT152: Food effect on the pharmacokinetics of 200-mg abemaciclib in healthy subject*. *Cancer Research*, 2016. **76**(14_Supplement): p. CT152-CT152.
9. Schmidt, L.E. and K. Dalhoff, *Food-drug interactions*. *Drugs*, 2002. **62**(10): p. 1481-502.
10. Chappell, J.C., et al., *Abemaciclib Inhibits Renal Tubular Secretion Without Changing Glomerular Filtration Rate*. *Clin Pharmacol Ther*, 2019. **105**(5): p. 1187-1195.
11. Schwartzberg, L., *Electronic Patient-Reported Outcomes: The Time Is Ripe for Integration Into Patient Care and Clinical Research*. *American Society of Clinical Oncology Educational Book*, 2016(36): p. e89-e96.
12. Rugo, H.S., et al., *The association of early toxicity and outcomes for patients treated with abemaciclib*. *Journal of Clinical Oncology*, 2018. **36**(15_suppl): p. 1053-1053.
13. Basch, E., et al., *Overall Survival Results of a Trial Assessing Patient-Reported Outcomes for Symptom Monitoring During Routine Cancer Treatment*. *Jama*, 2017. **318**(2): p. 197-198.
14. Basch, E., et al., *Symptom Monitoring With Patient-Reported Outcomes During Routine Cancer Treatment: A Randomized Controlled Trial*. *J Clin Oncol*, 2016. **34**(6): p. 557-65.

A. Grade 2 Diarrhea Events




B. Grade 3 Diarrhea Events



JPCP = current study. Abbreviations: M1 = MONARCH 1, M2 = MONARCH 2, M3 = MONARCH 3;


N= total number of patients in the abemaciclib treatment group.

Data cutoff: JPCP, 8 July 2019; MONARCH 1, 30 April 2016; MONARCH 2, 14 February 2017; MONARCH 3, 31 January 2017.



Abbreviations: N = number of patients assessed for eligibility and randomized, n = number of patients within category.

^aAt the time of data cutoff on 8 July 2019




Median = center line, 25th and 75th percentiles = hinges, and 1.5*IQR (inter-quartile range) = whiskers.
All individual data are represented on the plot as filled circles.

Fig 4A


[Click here to access/download;Figure;Fig 4A.eps](#)

Toxicity Grade
0 1 2 3
■ ■ ■ ■



Treatment durations represented are from first 3 cycles only but appear >84 days due to dose delays or dose interruptions

- Each row represents one patient
- Each square represents one day



*Number displayed in each bar is the number of patients with specific grade of diarrhea within each cycle.

Fig 4C

[Click here to access/download;Figure;Fig 4C.eps](#)






Fig 4D



*Number displayed in each bar is the number of patients with specific grade of diarrhea within each cycle.




Treatment Day


Treatment durations represented are from first 3 cycles only but appear >84 days due to dose delays or dose interruptions

- Each row represents one patient
- Each square represents one day
- Numbers represent numbers of loperamide 2mg tablets taken each day
- Number 8 assumed to represent number of abemaciclib tablets taken on that day (entered in error by patient)
- Patient-reported grade of diarrhea calculated from number of daily BM reported in e-diary with respect to baseline assessment



Click here to access/download
Supplementary Material
Supplementary Information.docx





[Click here to access/download](#)

Supplemental Materials - Not to be Published
CONSORT 2010 Checklist.doc

